ARRAY 2

Intestinal Antigenic Permeability Screen™

2 mL Serum

Antigens Tested

- Actomyosin IgA
- Occludin/Zonulin IgG
- Occludin/Zonulin IgA
- Occludin/Zonulin IgM
- Lipopolysaccharides (LPS) IgG
- Lipopolysaccharides (LPS) IgA
- Lipopolysaccharides (LPS) IgM

CLINICAL USE:	RECOMMENDED FOR PATIENTS WHO:
• Measure intestinal permeability to large molecules which inflame the immune system. • Identify the damaging route through the intestinal barrier.	• Present multiple symptom complaints (including Chronic Fatigue Syndrome). • Suffer from abnormal immune cell count and function (including autoimmune diseases). • Have multiple food sensitivity.

Cyrex® Laboratories

www.Joincyrex.com

©2017 Cyrex Laboratories, LLC. All Rights Reserved.

Transcellular & Paracellular Routes of Intestinal Barrier Penetration



Antigens Measured	
Actomyosin Proteins	
Occludin/Zonulin Proteins	
Lipopolysaccharides	Endotoxins from Gram-Negative Bacteria

Illustrations dramatized for visual purpose.
Image property of Cyrex Labs. © 2017 Cyrex Laboratories, LLC. All rights reserved.
A2BRO-05172017